



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/82, A01H 4/00	A1	(11) International Publication Number: WO 00/53783 (43) International Publication Date: 14 September 2000 (14.09.00)
(21) International Application Number: PCT/SG99/00016 (22) International Filing Date: 10 March 1999 (10.03.99) (71) Applicant (for all designated States except US): INSTITUTE OF MOLECULAR AGROBIOLOGY [SG/SG]; 1 Research Link, Singapore 117604 (SG). (72) Inventors; and (75) Inventors/Applicants (for US only): JIAO, Gai-Li [CN/SG]; Block 322, Clementi Avenue 5, #03-237, Singapore 120322 (SG). LIU, Jian, Wei [-/SG]; 13 Toh Yi Drive #07-05, Singapore 590013 (SG). (74) Agent: ELLA CHEONG & G. MIRANDAH; P.O. Box 0931, Raffles City, Singapore 911732 (SG).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: AGROBACTERIUM-MEDIATED TRANSFORMATION OF COTTON WITH NOVEL EXPLANTS (57) Abstract A method is disclosed for producing a transgenic cotton plant comprising the steps of (a) obtaining cottonfibrous root explants, (b) culturing the fibrous root explants to induce callus formation, (c) exposing root callus to a culture of <i>Agrobacterium tumefaciens</i> that harbors a vector comprising an exogenous gene and a selectable marker, the <i>Agrobacterium</i> being capable of effecting the stable transfer of the exogenous gene and selection agent resistance gene to the genome of the cells of the explant, (d) culturing the callus in the presence of the selection agent to which the selection agent resistance gene confers resistance so as to select for transformed cells, (e) inducing somatic embryo formation in the selected callus culture, and (f) regenerating the induced somatic embryos into whole transgenic cotton plants.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Agrobacterium-Mediated Transformation of
Cotton With Novel Explants

Technical Field

5 The present invention relates to the general field
of genetic engineering of plants, in particular to the
introduction of exogenous genetic material into cotton
by *Agrobacterium* transformation of novel explants
followed by somatic embryo regeneration.

Background

10 Cotton is the most extensively used natural fiber
in the textile industry. Its annual production
worldwide is over 100 million bales, valued at US\$45
billion. Cotton lint or seed hair is a terminally
differentiated single epidermal cell from 50 species of
15 the genus *Gossypium* of the family Malvaceae. It is
classified as a natural, cellulosic, monocellular and
staple fiber. The cultivated cotton varieties, which
have been cultivated for more than 5000 years, all come
from two diploids ($2n=2x=26$) (*G. herbaceum* and *G.*
20 *arboreum*) and two allotetraploids ($2n=4x=52$) (*G.*
hirsutum L., Upland; and *G. barbadense* L., Sea Island).
In 1997 the top five world cotton producers were the
United States, China, India, Pakistan and Uzbekistan,
producing about 63 million bales.

In the next century, most crops, including cereals, oil crops, fruits, vegetables and other economically important crops, will be genetically engineered with added or modified traits ranging from improvement of yield and quality, to herbicide resistance and pest resistance (Chappell, 1996; Fraley et al., 1986; Herrera-Estrella et al., 1983; Hoekema et al., 1983; Horsch et al., 1985; Jefferson, 1987; Ryals, 1996). In cotton, the new technology will be used to increase yield, improve fiber quality and create new varieties which are resistant to herbicides, pest insects, nematodes and diseases (John, 1996; John & Keller, 1996; John & Stewart, 1992; Murray et al., 1993; Rajasekaran et al., 1996; Schell, 1997; Stewart, 1992).

1. Tissue Culture of Cotton: In 1935, Skovsted reported the first embryo culture of cotton. Beasley (1971) reported callus formation in cotton as an outgrowth from the micropylar end of fertilized ovules on MS medium. Somatic embryogenesis was achieved from a suspension culture of *G. klotzschianum* (Prive & Smith, 1979). In 1983, Davidonis & Hamilton first succeeded in efficient and repeatable regeneration of cotton (*G. hirsutum* L.) plants from callus after two-year cultivation. Cotton plants were since regenerated through somatic embryogenesis from different explants (Zhang & Feng, 1992; Zhang, 1994) including cotyledon (Davidonis et al., 1987; Davidonis & Hamilton, 1983; Finer, 1988; Firoozabady et al., 1987), hypocotyl (Cousins et al., 1991; Rangan & Zavala, 1984; Rangan & Rajasekaran, 1996; Trolinder & Goodin, 1988; Umbeck et al., 1987, 1989), stem (Altman

et al., 1990; Finer & Smith, 1984), shoot apex (Bajaj et al., 1985; Gould et al., 1991; Turaev & Shamina, 1986), immature embryo (Beasley, 1971; Eid et al., 1973; Stewart & Hsu, 1977, 1978), petiole (Finer & Smith, 1984; Gawel et al., 1986; Gawel & Robacker, 1990), leaf (Finer & Smith, 1984; Gawel & Robacker, 1986), root (Chen & Xia, 1991; Kuo et al., 1989), callus (Finer & McMullen, 1986; Trolinder et al., 1991) and protoplast (Chen et al., 1989).

2. Cotton Transformation: Explants (such as hypocotyl, cotyledon, callus generated from hypocotyl and cotyledon, as well as immature embryos) have been used for *Agrobacterium*-mediated transformation and particle bombardment (de Framond et al., 1983; Finer & McMullen, 1990; Firoozabady et al., 1987; Perlak et al., 1990; Rangan & Rajasekaran, 1996; Rajasekaran et al., 1996; Trolinder et al., 1991; Umbeck et al., 1987, 1989, 1992). In addition, meristematic tissue of excised embryonic axes has also been used for cotton transformation by particle bombardment (Chlan et al., 1995; John, 1996; John & Keller, 1996; McCabe & Martinell, 1993). Zhou et al. (1983) transformed cotton by injecting DNA into the axile placenta one day after self-pollination. However, cotton transformation is highly dependent on genotype (Trolinder, 1985a, 1985b, 1986; Trolinder & Goodin, 1987, 1988a, 1988b). Apart from a few cultivars which are regeneratable and transformable, such as *Gossypium hirsutum* cv. Coker 312 and *G. hirsutum* Jin 7, most other important elite commercial cultivars, such as *G. hirsutum* cv. D&P 5415 and *G. hirsutum* cv. Zhongmian 12, are not regeneratable and transformable by these methods.

Based on previous reports and the inventor's own experimental data, high efficiency of callus induction (60%) can be achieved using the hypocotyl as an explant. However, the transformation rate was only 20% (Firoozabady et al., 1987; Umbeck et al., 1987). Several factors can lead to breakthrough of nontransformed calli, or to chimeric calli consisting of predominantly nontransformed cells: (1) low kanamycin levels (a high level of kanamycin is toxic to cotton explants and calli); (2) experience-dependent selection in later stages of callus proliferation; and (3) use of explants such as the hypocotyl which has only partial contact with the selective media (Firoozabady et al., 1987). When the cotyledon was used as an explant, although the transformation rate was higher than that with the hypocotyl, it was often difficult to eliminate *Agrobacterium* during subsequent culture (Jiao G.-L and Chen, Z.-X., personal communication; Umbeck et al., 1987, 1989). The transformation rate of meristemic tissue through particle bombardment was simply too low (0.02%-0.22%) compared to that of *Agrobacterium* mediated transformation.

There thus remains a need for methods of producing transgenic cotton plants that provide high rates of transformation along with high rates of transformants among regenerated somatic embryos.

Summary of the Invention

The present invention relates to a method for producing transgenic cotton plants, comprising the steps of (a) obtaining cotton fibrous root explants, (b) culturing the fibrous root explants to induce callus formation, (c) exposing root callus to a culture of *Agrobacterium tumefaciens* that harbors a vector comprising an exogenous gene and a selectable marker, the *Agrobacterium* being capable of effecting the stable transfer of the exogenous gene and selection agent resistance gene to the genome of the cells of the explant, (d) culturing the callus in the presence of the selection agent to which the selection agent resistance gene confers resistance so as to select for transformed cells, (e) inducing somatic embryo formation in the selected callus culture, and (f) regenerating the induced somatic embryos into whole transgenic cotton plants.

The present method provides for an improved rate of transformation when compared to previous methods that employ hypocotyl and cotyledon tissue. The method is believed to have wide applicability to a variety of cotton varieties.

Brief description of the Figures

Figure 1 shows the plasmid pBK9, containing a luciferase gene used to detect positive transformants obtained by the methods of the present invention.

Figure 2 shows the plasmid pVIP96, the plasmid from which pBK9 was derived by insertion of the luciferase gene.

Detailed Description

In order to overcome the problems seen with prior art methods and increase the efficiency of transformation, fibrous root explants were used for *Agrobacterium*-mediated transformation of cotton. Although in *Arabidopsis* high efficiency of transformation was achieved in *Agrobacterium*-mediated transformation with fibrous root explants (Valvekens, et al., 1988), and the differentiation of young fibrous roots from cotton on MS medium containing 2.0 mg/L IAA, 0.02-0.04 mg/L IBA has been reported (Kuo, C.C., et al., 1989), there is no report in the literature about using fibrous roots as explants for cotton transformation.

Fibrous roots now have been successfully used as explants for *Agrobacterium*-mediated transformation and plant regeneration. In the process modified media for seedling culture, and regeneration and differentiation of embryogenic calli were used.

Media used to culture seedlings to obtain explant material was designed to minimize browning of the roots (browning adversely effects the ability of explants to grow in culture and form callus), and to promote overall vigorous root growth. In a preferred embodiment MET (multi-effect triazole, a chemical agent used in agriculture to promote root growth) and NAA (α naphthalene acetic acid) are used together in the seedling culture medium to reduce the proportion of browned roots and increase callus initiation rate. MET is preferably used in concentrations ranging from about 0.05 mg/l to about 0.2 mg/l, most preferably about 0.1 mg/l. NAA is preferably used in concentrations ranging

from about 0.01 mg/l to about 0.2 mg/l, most preferably about 0.05 mg/l. MET and NAA are also preferably used in the medium used to root transgenic seedlings regenerated from callus, in amounts similar to those described for the seedling culture medium. In a preferred embodiment of the callus-forming medium vitamin B₅, 2,4-D ((2,4-dichlorophenoxy)acetic acid, MgCl and glucose are used, preferably about 0.05 mg/l to about 0.15 mg/l 2,4-D, about 0.4 mg/l to about 1.2 mg/l MgCl, and about 1% to about 5% glucose, most preferably about 0.1 mg/L 2,4-D, 0.8 mg/L MgCl and 3% glucose. In an alternate preferred embodiment of the callus-forming medium myo-inositol, vitamin B₁, and dimethylallyl(amino)purine are used, a, preferably about 50 mg/l to about 150 mg/l myo-inositol, about 1 mg/l to about 10 mg/l vitamin B₁, and about 0.1 mg/l to about 7.5 mg/l dimethylallyl(amino)purine, most preferably about 100 mg/l myo-inositol, about 0.4 mg/l vitamin B₁ and about 5 mg/l dimethylallyl(amino)purine. The same media used for callus induction can also be used during selection with antibiotics -- for example with 300-400 mg/L cefotaxime or 15-30 mg/L kanamycin. The presence of high concentrations (preferably about 1900 mg/l to about 5700 mg/l, most preferably about 3800 mg/L) of nitrates (preferably NaNO₃) was crucial for the observed effectiveness of the differentiation medium. With the fibrous roots as explants, although the rate of callus-induction was lower compared with hypocotyl and cotyledon, a higher rate of transformation was achieved.

Techniques for introducing exogenous genes into *Agrobacterium* such that they will be transferred stably

to a plant or plant tissue exposed to the *Agrobacterium* are well-known in the art and do not form part of the present invention. It is advantageous to use a so-called "disarmed" strain of *Agrobacterium* or Ti

5 plasmid, that is, a strain or plasmid wherein the genes responsible for the formation of the tumor characteristic of the crown gall disease caused by wild-type *Agrobacterium* are removed or deactivated. Numerous examples of disarmed *Agrobacterium* strains are

10 found in the literature (e.g., pAL4404, pEHA101 and pEH105 (Walkerpeach & Veltner, 1994)). It is further advantageous to use a so-called binary vector system, such as that described in U.S. Patent Nos. 4,940,838 and 5,464,763 (Schilperoort, et al.) and Hoekema et

15 al., 1983. A binary vector system allows for manipulation in *E. coli* of the plasmid carrying the exogenous gene to be introduced into the plant, making the process of vector construction much easier to carry out.

20 Similarly, vector construction, including the construction of chimeric genes comprising the exogenous gene that one desires to introduce into the plant, can be carried out using techniques well-known in the art and does not form part of the present invention.

25 Chimeric genes should comprise promoters that have activity in the host in which expression is desired. For example, it is advantageous to have a series of selectable markers for selection of transformed cells at various stages in the transformation process. A

30 selectable marker (for example a gene conferring resistance to an antibiotic such as kanamycin, cefotaxime or streptomycin) linked to a promoter active

in bacteria would permit selection of bacteria containing the marker (i.e., transformants). Another selectable marker linked to a plant-active promoter, such as the CaMV 35S promoter or a T-DNA promoter such as the NPT II NOS promoter, would allow selection of transformed plant cells. The exogenous gene that is desired to be introduced into the plant cell should comprise a plant-active promoter in functional relation to the coding sequence, so that the promoter drives expression of the gene in the transformed plant. Again, plant-active promoters, such as the CaMV 35S, the NPT II NOS promoter or any of a number of tissue-specific promoters, are well-known in the art and selection of an appropriate promoter is well within the ordinary skill in the art.

The present method can be used to produce transgenic plants expressing any number of exogenous genes, and is not limited by the choice of such a gene. The selection of the desired exogenous gene depends on the goal of the researcher, and numerous examples of desirable genes that could be used with the present invention are known in the art (e.g., the family of *Bacillus thuringiensis* toxin genes, herbicide resistance genes such as shikimate synthase genes that confer glyphosate resistance, U.S. Patent No. 5,188,642, or a 2,4-D monooxygenase gene that confers 2,4-D resistance, Bayley et al., Theoretical and Applied Genetics, vol. 82, pp. 645-49, male sterility genes such as the antisense genes of U.S. Patent No. 5,741,684 (Fabijanski, et al.), or even the elaborate crop protection systems described in U.S. Patent No. 5,123,765 (Oliver, et al.)).

Agrobacterium-mediated cotton transformation is considered in the art to be heavily variety-dependant. The Coker series of cotton varieties have been shown to be relatively easy to transform. However, DP 5412, 5 Zhongmain 12 and many other varieties still have difficulties associated with transformation. The situation is the same for *G. barbadense* and other diploid species. Particle bombardment, DNA injection and infection of meristem tissue with *Agrobacterium* are 10 some alternative methods, which can be used to transform, in theory, all the cotton varieties. The problems associated with these methods are: low efficiency of transformation and unstable/unreliable results. It is believed that the present method has 15 broad applicability to transformation of cotton varieties, as it overcomes or minimizes several of the problems associated with previous work relating to cotton transformation (such as breakthrough of non-transformed callus, poor explant growth and low 20 transformation rate, poor somatic regeneration) through the use of fibrous root explants.

The following abbreviations are used to designate culture media useful in connection with the present invention:

25 LB medium (10g bacto-tryptone + 5g bacto-yeast extract + 10g NaCl);
 B5 medium (Gamborg et al., 1968; Sigma, Cat. No. G-5768);
 MS medium (Murashige et al., 1962; Sigma, Cat. No. 30 M-5524);
 SH medium (Stewart & Hsu, 1977. Planta 137, 113-117);

- CB-1.1 ($\frac{1}{2}$ MS + $\frac{1}{2}$ B5 Vitamin + 0.1mg/L NAA);
CB-1.2 ($\frac{1}{2}$ B5 medium);
CB-2.1 (MS macro + B5 micro + 0.05mg/L 2,4-D +
0.1mg/L kinetin + 3% glucose + 2g/L gellan gum
(PhytaGel™, Sigma) + 0.93mg/L MgCl₂·6H₂O, pH5.8);
CB-2.2 (MS macro + 100mg/L myo-inositol + 0.4mg/L
vitamin B1 + 5mg/L 2iP (6 - (γγ-
dimethylallyl(amino)purine)+ 0.2mg/L NAA + 3%
glucose + 2g/L gellan gum (PhytaGel™, Sigma) +
0.93mg/L MgCl₂·6H₂O, pH5.8);
CB-3.1 (CB-2.1+ 500 mg/L cefotaxime + 50mg/L
kanamycin);
CB-3.2 (CB-2.2 + 500mg/L cefotaxime + 50mg/L
kanamycin);
CB-4 (Modified CB-3.1 or CB-3.2 by adding double
amount of KNO₃ and removing NH₄NO₃ with 250 mg/L
cefotaxime and 20mg/L kanamycin);
CB-5 (SH + 1.5% Sucrose + 2g/L gellan gum
(PhytaGel™, Sigma) + 0.93g/L MgCl₂, pH7.0).

The following Examples are intended to illustrate the present invention, and not in any way to limit its scope, which is solely defined by the claims.

EXAMPLE 1: Regeneration of Cotton Plants from Root Tissue Culture

Preparation of Root Explants: Cotton seeds were sterilized in 70% ethanol for 10-15 min., and then treated with 10% H₂O₂ for 30-120 mins. Treated seeds were rinsed in sterile water for 24 hrs at 28°C and germinated on either CB-1.1 medium or CB-1.2 medium at 28°C - 30°C, 16h light (60-90 μE m⁻² s⁻¹). Seven to ten

days sterile seedlings thus grown were used to prepare explants. It was found that plentiful healthy roots (longer and thicker) with white color were obtained using CB-1.1 medium, whereas shorter and thinner roots with grey to brown color were obtained using the CB-1.2 medium. Therefor, CB-1.1 was chosen for further work.

Induction of calli: Fibrous roots were cut from seedlings and cultured on CB-2.1 medium or CB-2.2 medium at 28°C - 30°C for three days, 16h light (60-90 $\mu\text{E m}^{-2} \text{s}^{-1}$). The optimum size for root explants was 5-7 mm. A few small calli initiated on the cut sites of root segments in as little as 3 days. In general, transformed hypocotyl or cotyledon explants started to initiate callus on inducing medium after 3 days. However, previous to the present invention, transformed root explants were generally found to initiate callus only after 10 days of cultivation. The color of the root explants was white. One week later, small calli were also initiated from other parts of the root segments. The color of the root explants changed to grey or even brown. At the end of 2 weeks of cultivation, calli initiated from the whole root explants and grew well. Of the two inducing media, CB-2.2 was found to induce good callus formation, while CB-2.1 did not. On CB-2.2 medium, root explants grew well, and the microcallus initiated on the cut sites of the explant. About 10% of root explants initiated callus after only 3 days on the medium. On the other hand, while CB-2.1 medium supported the growth of the root explants well, there was no callus initiation on the cut sites. The efficiency of callus-induction with root explants on the CB-2.2 medium was 10%, which was

lower than that with hypocotyl or cotyledon explants (20-30%). A summary of results showing callus induction and transformation efficiency appears in Table 1, below.

5 Regeneration of root calli: After one month, the calli were transferred to new medium for subculture on either CB-2.1 medium or CB-2.2 medium. After 2 months of subculture, the mature calli were transferred to CB-4 (without antibiotic) for induction of somatic
10 embryos. Glutamine and L-asparagine were added in amounts of 0.5 mg/L and 0.2 mg/L, respectively, to promote embryogenesis. Primary somatic embryos were formed on the embryogenic calli after 2 months of cultivation, with 2 subcultures in between on the same
15 media. Primary somatic embryos were subcultured on the same media for another month before mature somatic embryos were formed. Some of the somatic embryos developed to plantlets. These small plantlets were transferred to CB-5 medium for root induction. When
20 the plantlets had made roots on the CB-5 medium (4-6 weeks), they were transferred to soil and maintained in an incubator under high humidity for 3-4 weeks at 28°C, 16h light (60-90 $\mu\text{E m}^{-2} \text{ s}^{-1}$), and then transferred to large pots with soil in a green house.

25 EXAMPLE 2: *Agrobacterium* Transformation and Culture

 The plasmid pBK9 (35S:LUC) (see Fig. 1) was generated by cloning the luc coding sequence from the BamHI/StuI fragment of plasmid pGEMluc into the blunt-ended StuI site of the plasmid pVIP96 (see Fig.
30 2).

Prepared competent cells (400 microliter) in Eppendorf tube from -80°C were put on ice to thaw. Plasmid DNA was added in the cells. After gentle mixing, the mixture was incubated on ice for 45 minutes. The Eppendorf tube containing the mixture was put into liquid nitrogen for 1 minute and afterwards in a water bath (37°C) for 3 minutes. After the incubation, 800 microliter LB medium (without antibiotics) was added into the mixture and the tube with the mixture was incubated at 28°C for 3 hours. After a brief centrifugation at 12,000 rpm, 800 microliter supernatant was removed. The rest of the medium was mixed well with the cell pellet and the mixture was plated onto LB plates containing 100 mg/L kanamycin and 100 mg/L streptomycin. Successful transformed LBA4404 cells formed colonies on the plates in about 48 hours at 28°C. *Agrobacterium* strain LBA4404 harboring the plasmid pBK9 (35S:LUC) was initiated on LB plate with kanamycin (50 mg/L), streptomycin (50 mg/L) and rifampicin (50 mg/L). A single colony was inoculated into LB liquid medium without antibiotics and grown overnight for about 18 h at 28°C on a gyratory shaker. The optical density (A_{600}) value was adjusted to 0.1 - 0.4 in liquid LB medium prior to use.

EXAMPLE 3: Transformation of Root Explant Tissue and Regeneration of Transgenic Cotton Plants

Root explants were obtained by cultivating sterile cotton seeds as described in Example 1, above, on CB-1.1 medium. Fibrous roots were cut from seedlings and

cultured on CB-2.2 for two days, 16h light ($60-90 \mu\text{E m}^{-2}\text{s}^{-1}$). The fibrous roots were then cut into small segments (5-10 mm) and incubated with the cell suspension culture of *Agrobacterium tumefaciens* strain LBA4404 harboring the plasmid pBK9 (35S:LUC) ($A_{600} = 0.1-0.6$) of Example 2 for 15 min. After drainage of the bacterial solution, the root explants were cultured at 28°C , 16h light ($60-90 \mu\text{E m}^{-2}\text{s}^{-1}$) for an additional two days. The optimum concentration of the *Agrobacterium* strain LBA4404 for root explants was lower ($A_{600}=0.1-0.4$) than that for hypocotyl and cotyledon explants ($A_{600}=0.3-0.6$). Optimal bacterial concentrations did not affect the growth of the root explants and the subsequent callus induction.

Co-cultured explants were washed twice with sterile distilled water and transferred to CB-3.1 medium or CB-3.2 medium for cultivation at 28°C , 16h light ($60-90 \mu\text{E m}^{-2} \text{s}^{-1}$).

After four weeks, kanamycin-resistant calli were selected and subcultured on the same media for the second selection. At the same time, some of the calli were selected to detect the LUC expression with the luciferase luminescence image system (see Example 4, below). The process of inducing callus took about 2 months. The efficiency of callus induction from root explant was lower compared with that from hypocotyl and cotyledon explants.

Kanamycin-resistant calli were transferred to CB-4 medium to induce embryogenic calli and somatic embryos. After 4-6 weeks of cultivation, with one subculture, mature somatic embryos appeared on the calli. Plantlets developed afterwards from some of the

embryos. The green plantlets were then transferred to rooting medium (CB-5) for root induction. When plantlets had made roots, they were transferred to soil and maintained in an incubator under high humidity for 3-4 weeks at 28°C, 16h light ($60-90 \mu\text{E m}^{-2} \text{s}^{-1}$), and then transferred to large pots with soil in a green house.

EXAMPLE 4: Detection of Luciferase Activity

Plant materials (such as callus, leaf and whole plantlet) were sprayed with a solution containing 0.5 mM potassium luciferin and 0.01% (w/v) polyoxyethylenesorbitan monolaurate (Tween-20) and left for 30 min. The luciferase luminescence from these plant materials was visualized using an image-intensifying camera and photon-counting image processors purchased from Princeton Instruments Inc., 3660 Quakerbridge Road, Trenton, NJ 08619. The exposure time was 6 min. The electronic images were converted to Microsoft Powerpoint TIFF files and printed out from a standard color printer.

Callus growing on selected medium for one month was selected to test LUC expression with the video image system. The positive transformed callus had white spots whereas untransformed callus did not. Out of the 139 pieces of kanamycin resistant calli, 49 pieces were positive with LUC activity. The successful transformation rate was therefor 35%, which was much higher than that seen using cotyledon or hypocotyl as explant (20%).

Table 1. Brief summary of root as explant for transformation

Variety	Construct	Conc. of Strain	Date of selected medium	No. of explant	No. of callus	No. of LUC test	No. of LUC+
5 Coker 312	35S-LUC	0.1-0.2	9/10/97	2574	156	16	12
	35S-LUC	0.43	23/10/97	1255	70	8	5
	35S-LUC	0.438	23/10/97	564	46	28	19
	35S-LUC	0.2-0.4	5/3/98	520	33	15	11

References

- Altman, D. W. et al. 1990. Economic Botany 40, 106.
- Bajaj, Y. P. S. 1985. Theor. Appl. Genet. 70, 363.
- 10 Beasley, C. A. 1971. In vitro culture of fertilized cotton ovules. Biosci. 21, 906-7.
- Chappell, J. 1996. Plant biotechnology comes of age - again. Mol. Breeding 2, 1-6.
- Chen, Z. X., Li, S. J., Yue, J. X., Jiao, G. L. and Liu S. X. 1989. Plantlet regeneration from protoplasts
15 isolated from an embryogenic suspension culture of cotton (*Gossypium hirsutum* L.). Acta Botanica Sinica 31, 966-9.
- Chen, Z. X. and Xia Z. A. 1991. Acta Botanica Sinica 33, 98.
- 20 Chlan, C. A., Lin, J., Cary, J. W. and Cleveland, T. E. 1995. A procedure for biolistic transformation and regeneration of transgenic cotton from meristematic tissue. Plant Mol. Biol. Rep. 13, 31-7.
- 25 Cousins, Y.L., Lyon, B. R. and Llewellyn, D. J. 1991.

- Transformation of an Australian cotton cultivar:
prospects for cotton improvement through genetic
engineering. Aust. J. Plant Physiol. 18, 481-94.
- Davidonis, G. H., and Hamilton, R. H. 1983. Plant
5 regeneration from callus tissue of *Gossypium*
hirsutum L. Plant Sci. Lett. 32, 89-93.
- Davidonis, G. H., Mumma, R. O. and Hamilton, R. H.
1987. Controlled regeneration of cotton plants
from tissue culture. US Patent No. 4,672,035.
- 10 de Framond et al. 1983. Mini-Ti: a new vector strategy
for plant genetic engineering. Bio/technology 1,
262-9.
- Finer, J. J. and Smith, R. H. 1984. Initiation of
callus and somatic embryos from explants of mature
15 cotton (*Gossypium klotzschianum* Anderss). Plant
Cell Reports 3, 41-43.
- Finer, J., 1988. Plant regeneration from somatic
embryogenic suspension cultures of cotton
(*Gossypium hirsutum* L.). Plant Cell Rep. 7,
20 399-402.
- Finer, J. J. and McMullen, M. D. 1990. Transformation
of cotton (*Gossypium hirsutum* L.) via particle
bombardment. Plant Cell Reports 8, 586-9.
- Firoozabady, E., Deboer, D., Merlo, D., Halk, E.,
25 Amerson, L., Rashka, K. and Murray, E. 1987.
Transformation of cotton (*Gossypium hirsutum* L.)
by *Agrobacterium tumefaciens* and regeneration of
transgenic plants. Plant Mol. Biol. 10, 105-16.
- Fraley, R. T., Rogers, S. G. and Horsch, R. B. 1986.
30 Genetic transformation in higher plants. CRC Cri.
Rev. Plant Sci. 4, 1-46.
- Gamborg, O., Miller, R. and Ojima, K. 1968. Exp. Cell

- Res. 50, 148.
- Gawel, N. J., Rao, A. P. and Robacker, C. 1986.
Somatic embryogenesis from leaf and petiole callus
cultures of *Gossypium hirsutum* L. Plant Cell
5 Reports 5, 4579.
- Gawel, N. J. and Robacker, C. 1986. Plant Cell Rep. 5,
67.
- Gawel, N. J. and Robacker, C. 1990. Genetic control of
somatic embryogenesis in cotton petiole callus
10 cultures. Euphytica 49, 249-53.
- Gould, J., Banister, S., Hasegawa, O., Fahima, M. and
Smith, R. H. 1991. Regeneration of *Gossypium*
hirsutum and *G. barbadense* from shoot apex tissues
for transformation. Plant Cell Reports 10, 12-6.
- 15 Herrera-Estrella, L., Depicker, A., Van Montagu, M. and
Schell, J. 1983. Expression of chimeric genes
transferred into plant cells using a
Ti-plasmid-derived vector. Nature 303, 209-13.
- Hoekema et al. 1983. A binary plant vector strategy
20 based on separation of Vir- and T-Region of the
Agrobacterium tumefaciens Ti-plasmid. Nature 303,
179-80.
- Horsch, R. B., Fry, J. E., Hoffmann, N. L., Eichholtz,
D., Rogers, S. G. and Fraley, R. T. 1985. A
25 simple and general method for transferring genes
into plants. Sci. 227, 1229-31.
- Jefferson, R. A. 1987. Assaying chimeric genes in
plants: the GUS gene fusion system. Plant Mol.
Biol. Rep. 5, 387-405.
- 30 John, M. E. 1996. Structural characterization of genes

- corresponding to cotton fiber mRNA, E6: reduced E6 protein in transgenic plants by antisense gene. Plant Mol. Biol. 30, 297-306.
- John, M. E. and Keller, G. 1996. Metabolic pathway engineering in cotton: Biosynthesis of polyhydroxybutyrate in fiber cells. Proc. Natl. Acad. Sci. USA 93, 12768-73.
- John, M. E., and Stewart, J. McD. 1992. Genes for jeans: biotechnological advances in cotton. Trends Biotechnol. 10, 165070.
- Kuo, C. C. et al., 1989. Proc. Beltwide Cotton Prod. Res. Confs, 638.
- McCabe, D. E. and Martinell, B. J. 1993. Transformation of elite cotton cultivars via particle bombardment of meristems. Bio/technol. 11, 596-8.
- Murray, E. E., DeBoer, D. L. and Firoozabady, E. 1993. Transgenic cotton. In "Transgenic Cotton", Vol 2, 153-68.
- Murashige et al., 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. Physiologia Plantarum 15, 473-97.
- Perlak, F. J., Deaton, R. W., Armstrong, T. A., Fuchs, T. A., Sims, S. R., Greenplate, J. T. and Fischhoff, D. A. 1990. Insect resistant cotton plants. Bio/technol. 8, 939-943.
- Price, H. J. and Smith, R. H. 1979. Somatic embryogenesis in suspension cultures of *Gossypium klotzschianum* Anderss. Planta 145, 305-6.
- Rajasekaran, K., Grula, J. W., Hudspeth, R. L., Pofelis, S. and Anderson, D. M. 1996. Herbicide-resistant Acala and Coker cottons

- transformed with a native gene encoding mutant forms of acetohydroxyacid synthase. *Mol. Breeding* 2, 307-19.
- 5 Rangan, F. J. and Zavala, T. Ip. A. 1984. Somatic embryogenesis in tissue culture of *Gossypium hirsutum* L.). *In Vitro* 20, 256.
- Rangan, R. and Rajasekaran, K. 1996. Regeneration of cotton plant in cell suspension culture. US Patent No. 5,583,036 (continued from US Patent 10 5,244,802, 1993 and 122,200, 1987).
- Ryals, J. 1996. *Agricultural Biotechnology '96*. *Molecular Breeding* 2, 91-93.
- Schell, J. 1997. Cotton carrying the recombinant insect poison Bt toxin: no case to double the benefits of 15 plant biotechnology.
- Skovsted, A. 1935. Cytological studies in cotton. III. A hybrid between *Gossypium davidsonii* Kell, and *G. sturrtii* F. Muell. *J. Genet.* 30, 397-405.
- 20 Stewart, J. McD. 1992. *Biotechnology of cotton: achievements and perspectives*. ICAC Review Articles on Cotton Production Research No. 3, 1-50.
- 25 Stewart, J. McD and Hsu, C. L. 1978. Hybridization of deploid and tetraploid cottons through in-ovulo embryo culture. *J. Heridity* 69, 404-8.
- Srewart, J. McD. and Hsu, C. L., 1977. In ovulo embryo culture and seedling development of cotton (*Gossypium hirsutum* L.). *Planta* 137, 113-7.
- 30 Thomas, J. C., Adams, D. G., Keppene, V. D., Wasmann, C. C., Brown, J. K., Kanost, M. R. and Bohnert, H. J. 1995. Protease inhibitors of *Manduca sexta*

- expressed in transgenic cotton. Plant Cell, Rep. 14, 758-62.
- 5 Trolinder, N. L. 1985a. Somatic embryogenesis and plant regeneration in cotton (*Gossypium hirsutum* L.). A Dissertation in Biology (Dec., 1985).
- Trolinder, N. L. 1985b. Genotype specificity of the somatic embryogenesis response in cotton. Plant Cell Reports 8, 133-6.
- 10 Trolinder, N. L. 1986. Somatic embryogenesis and plant regeneration in *Gossypium hirsutum* L. Dissertations Abstracts International 47, 2550B-2551B.
- 15 Trolinder, N. L. and Goodin, J. E. 1987. Somatic embryogenesis and plant regeneration in cotton (*Gossypium hirsutum* L.). Plant Cell Reports 6, 231-4.
- 20 Trolinder, N. L. and Goodin, J. E. 1988a. Somatic embryogenesis and regeneration in cotton. I. Effects of source of explant and hormone regime. Plant Cell Tissue Organ Culture 12, 31-42.
- Trolinder, N. L. and Goodin, J. E. 1988b. Somatic embryogenesis and regeneration in cotton. II. Requirements for embryo development and plant regeneration. Plant Cell Tissue Organ Culture 12, 25 43-53.
- Trolinder, N. L., Quisenberry, J., Bayley, C., Ray, C., and Ow, D. 1991. 2,4-D resistant transgenic cotton. Proc. Beltwide Cotton Prod. Res. Conf., 840.
- 30 Turaev, A. M. and Shamina, Z. B. 1986. Soniet Plant Physiol. 33, 439.
- Umbeck, P. F., 1991. Genetic engineering of cotton

- plants and lines. US Patent No. 5,004,863.
- Umbeck, P. F. 1992. Genetic engineering of cotton plants and lines. US Patent No. 5,159,135 (continued from US Patent 5,004,863, 1991).
- 5 Umbeck, P. F., Johnson, G., Barton, K. and Swain, W. 1987. Genetically transformed cotton (*Gossypium hirsutum* L.) plants. *Bio/technol.* 5, 263-6.
- Umbeck, P. F., Swain, W. and Yang, N.-S., 1989. Inheritance and expression of genes for kanamycin and chloramphenical resistance in transgenic
10 cotton plants. *Crop. Sci.* 29, 196-201.
- Valvekens, D., Van Montagu, M., Van Lijsebettens, M. 1988. *Agrobacterium tumefaciens*-mediated transformation of *Arabidopsis* root explants by using kanamycin selection. *Proc. Natl. Acad. Sci.*
15 USA 85, 5536-40.
- Walkerpeach, C.R. and Veltern, J. 1994. *Agrobacterium*-mediated gene transfer to plant cells: cointegrate and binary vector systems. *Plant Mol. Biol.*
20 *Mannuel B1*, 1-19.
- Zhang, H.-B. 1994. The tissue culture of cotton (II). *Plant Physiol. Commun.* 30, 386-91.
- Zhang, H.-B. and Feng, R. 1992. The tissue culture of cotton (I). *Plant Physiol. Commun.* 30, 386-91.
- 25 Zhou, G.-Y., Weng, J., Zeng, Y.-S., Huang, J.-G., Qian, S.-Y. and Liu, G.-L. 1983. Introduction of exogenous DNA into cotton embryos. *Methods in Enzymology* 101,433-81.

We claim:

1. A method for producing a transgenic cotton plant comprising the steps of:
 - (a) obtaining cotton fibrous root explants,
 - (b) culturing the fibrous root explants to
5 induce callus formation,
 - (c) exposing root callus to a culture of *Agrobacterium tumefaciens* that harbors a vector comprising an exogenous gene and a selectable marker, the *Agrobacterium* being
10 capable of effecting the stable transfer of the exogenous gene and selection agent resistance gene to the genome of the cells of the callus,
 - (d) culturing the callus in the presence of
15 the selection agent to which the selection agent resistance gene confers resistance so as to select for transformed cells,
 - (e) inducing somatic embryo formation in the selected callus culture, and
20 (f) regenerating the induced somatic embryos into whole transgenic cotton plants.
2. The method of claim 1 wherein the cotton fibrous root explants are obtained by growing cotton seedlings in the presence of multi-effect
25 triazole.
3. The method of claim 2 wherein the multi-effect triazole is in a concentration of about 0.05 mg/l to about 0.2 mg/l.

4. The method of claim 3 wherein the multi-effect triazole is in a concentration of about 0.1 mg/l.
5. The method of claim 2 wherein the cotton seedlings are grown in the additional presence of α naphthalene acetic acid.
5
6. The method of claim 5 wherein the α naphthalene acetic acid is in a concentration of about 0.01 mg/l to about 0.2 mg/l.
7. The method of claim 6 wherein the α naphthalene acetic acid is in a concentration of about 0.05 mg/l.
10
8. The method of claim 1 wherein the step of regenerating the somatic embryos is carried out in the presence of multi-effect triazole.
9. The method of claim 8 wherein the multi-effect triazole is in a concentration of about 0.05 mg/l to about 0.2 mg/l.
15
10. The method of claim 9 wherein the multi-effect triazole is in a concentration of about 0.1 mg/l.
11. The method of claim 8 wherein the step of regenerating the somatic embryos is carried out in the additional presence of α naphthalene acetic acid.
20

12. The method of claim 11 wherein the α naphthalene acetic acid is in a concentration of about 0.01 mg/l to about 0.2 mg/l.
- 5 13. The method of claim 12 wherein the α naphthalene acetic acid is in a concentration of about 0.05 mg/l.
- 10 14. The method of claim 1 wherein the step of inducing callus formation is carried out in a callus inducing culture medium comprising myo-inositol, vitamin B₁ and a dimethylallyl(amino)purine.
- 5 15. The method of claim 1 wherein the step of inducing somatic embryo formation is carried out in a somatic embryo inducing culture medium comprising myo-inositol, vitamin B₁, and a dimethylallyl(amino)purine.
- 5 16. The method of claim 14 wherein the callus inducing culture medium comprises myo-inositol in an amount from 50 mg/L to 150 mg/L, vitamin B₁ in an amount from 0.2 to 10 mg/L and a dimethylallyl(amino)purine in an amount from 0.1 to 7.5 mg/L.
17. The method of claim 16 wherein the callus inducing culture medium comprises 100 mg/L myo-inositol, 0.4 mg/L vitamin B₁ and 5 mg/L dimethylallyl(amino)purine.

18. The method of claim 15 wherein the somatic embryo inducing culture medium comprises myo-inositol in an amount from 50 to 100 mg/L, vitamin B₁ in an amount from 0.2 to 10 mg/L and
5 dimethylallyl(amino)purine in an amount from 0.01 to 0.5 mg/L.
19. The method of claim 18 wherein the somatic embryo inducing medium comprises 100 mg/L myo-inositol, 0.4 mg/L vitamin B₁ and 5 mg/L dimethylallyl(amino)purine.
20. The method of claim 1 wherein the step of inducing callus formation is carried out in a callus inducing culture medium comprising vitamin B₅, (2,4-dichlorophenoxy)acetic acid, MgCl and glucose.
21. The method of claim 1 wherein the step of inducing somatic embryo formation is carried out in a somatic embryo inducing culture medium comprising vitamin B₅, (2,4-dichlorophenoxy)acetic acid, MgCl
5 and glucose.
22. The method of claim 20 wherein the callus inducing culture medium comprises vitamin B₅ in an amount from 0.2 mg/L to 10 mg/L, (2,4-dichlorophenoxy)acetic acid in an amount from 0.05
5 mg/L to 0.15 mg/L, MgCl in an amount from 0.4 mg/L to 1.2 mg/L and glucose in an amount from 1% to 5%.

23. The method of claim 22 wherein the callus inducing culture medium comprises 0.4 mg/L vitamin B₅, 0.1 mg/L (2,4-dichlorophenoxy)acetic acid, 0.8 mg/L MgCl and 3% glucose.
24. The method of claim 21 wherein the somatic embryo inducing culture medium comprises vitamin B₅ in an amount from 0.2 mg/L to 10 mg/L, (2,4-dichlorophenoxy)acetic acid in an amount from 0.05 mg/L to 0.15 mg/L, MgCl in an amount from 0.4 mg/L to 1.2 mg/L and glucose in an amount from 1% to 5%.
25. The method of claim 24 wherein the somatic embryo inducing medium comprises 0.4 mg/L vitamin B₅, 0.1 mg/L (2,4-dichlorophenoxy)acetic acid, 0.8 mg/L MgCl and 3% glucose.
26. A method according to any of claims 14-25, wherein the medium further comprises gellan gum.
27. A method according to claim 26 wherein the gellan gum is present in an amount from 1.0 g/L to 3.0 g/L.
28. The method of claim 1 wherein the step of inducing somatic embryo culture is carried out in a somatic embryo-inducing medium comprising a nitrate in an amount from 1900 mg/L to 5700 mg/L.
29. The method of claim 28 wherein the somatic embryo-inducing medium comprises 3800 mg/L nitrate.

30. A method according to either claim 28 or 29,
wherein the nitrate is NaNO_3 .

1/2

FIG. 1





FIG. 2



INTERNATIONAL SEARCH REPORT

International Application No

PCT/SG 99/00016

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12N15/82 A01H4/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	DATABASE AGRICOLA 'Online! AN 92:20667 DN IND92003356, KUO, C.C., ET AL.: "Plant regeneration from root-derived callus of Gossypium hirsutum L" XP002126090 abstract & PROCEEDINGS - BELTWIDE COTTON CONFERENCES, 1989. (BOOK 2) P. 638-640. PUBLISHER: MEMPHIS, TENN.: NATIONAL COTTON COUNCIL OF AMERICA.,	1-7, 14-30
Y	WO 97 12512 A (CALGENE INC) 10 April 1997 (1997-04-10) page 11, line 7 - line 24 page 14, line 12 - page 15, line 2 -/-	1-7, 14-30

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "Z" document member of the same patent family

Date of the actual completion of the international search

17 December 1999

Date of mailing of the international search report

11/01/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Maddox, A

INTERNATIONAL SEARCH REPORT

Inte. onal Application No

PCT/SG 99/00016

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>DATABASE CAB 'Online! CAB INTERNATIONAL, WALLINGFORD, OXON, GB AN 96:45438, CHEN, S., ET AL.: "Studies on the physiological effect of ABT No.4 on cotton" XP002126091 abstract & ACTA AGRICULTURAE SHANGHAI, vol. 10, no. 1, 1994, pages 73-77,</p>	2-7
A	<p>WO 98 15622 A (DEVER JANE GAY KVETON ;TROLINDER NORMA L (US); KOONCE LINDA KAY TR) 16 April 1998 (1998-04-16) page 6, line 9 - line 39 page 12, line 27 - line 38; examples 2,3 page 10, line 21 - line 30</p>	1-30
A	<p>DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US AN 1991:426929, LU, Z-X, ET AL.: "Study on tissue and protoplast culture of wild cotton Gossypium-davidsonii" XP002126111 abstract & ACTA BOT SIN, vol. 33, no. 2, 1991, pages 98-103,</p>	1-30
A	<p>FIROOZABADY ET AL: "Transformation of cotton (Gossypium hirsutum L.) by Agrobacterium tumefaciens and regeneration of transgenic plants" PLANT MOLECULAR BIOLOGY, NL, NIJHOFF PUBLISHERS, DORDRECHT, vol. 10, 1997, page 105-116 XP002094283 ISSN: 0167-4412 page 107 -page 108</p>	14-19
A	<p>TROLINDER N L ET AL: "SOMATIC EMBRYOGENESIS AND PLANT REGENERATION IN COTTON (GOSSYPIMUM HIRSUTUM L.)" PLANT CELL REPORTS, DE, SPRINGER VERLAG, vol. 6, no. 3, 1987, page 231-234 XP000618864 ISSN: 0721-7714 the whole document</p>	20-25
A	<p>EP 0 270 355 A (AGRACETUS) 8 June 1988 (1988-06-08) the whole document</p>	20-25
	-/-	

INTERNATIONAL SEARCH REPORT

Int. l. Application No

PCT/SG 99/00016

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>DATABASE CAB 'Online! CAB INTERNATIONAL, WALLINGFORD, OXON, GB AN 1998:72033, HUI, L., ET AL.: "Studies on the application of multi-effect-triazole in anther culture in wheat" XP002126092 abstract & ACTA AGRONOMICA SINICA, vol. 23, no. 2, 1997, pages 220-225,</p>	2-13
A	<p>DATABASE CHEMABS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US AN 117:128291, SHANKLA, A., ET AL.: "Influence of growth regulators on somatic embryogenesis, plantlet regeneration, and post-transplant survival of Echinochloa frumentacea" XP002126129 abstract & PLANT CELL REP. (1992), 11(7), 368-71,</p>	8-13
A	<p>DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, USq AN 1991:22373, ZHAO, C-Z., ET AL.: "Effect of MET on immature embryo callus induction differentiation and hardening of plants" XP002126130 abstract & ACTA BOT SIN, vol. 32, no. 5, 1990, pages 407-409,</p>	8-13
A	<p>DATABASE CAB 'Online! CAB INTERNATIONAL, WALLINGFORD, OXON, GB AN 95:112675, PAN, X.Y., ET AL.: "A study on the effect of MET on regulating the growth of plantlet from rice pollen" XP002126093 abstract & ACTA AGRICULTURAE UNIVERSITATIS JIANGXIENSIS, vol. 15, no. 3, 1993, pages 225-229,</p>	2-13
A	<p>EP 0 317 511 A (CIBA GEIGY AG) 24 May 1989 (1989-05-24) page 31 -page 37</p>	14-30
A	<p>WO 89 05344 A (PHYTOGEN) 15 June 1989 (1989-06-15) the whole document</p>	14-30
	-/-	

INTERNATIONAL SEARCH REPORT

Int. l. Application No

PCT/SG 99/00016

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 4 672 035 A (DAVIDONIS GAYLE H ET AL) 9 June 1987 (1987-06-09) cited in the application the whole document	28-30
A	VALVEKENS D ET AL: "AGROBACTERIUM TUMEFACIENS-MEDIATED TRANSFORMATION OF ARABIDOPSIS THALIANA ROOT EXPLANTS BY USING KANAMYCIN SELECTION" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, US, NATIONAL ACADEMY OF SCIENCE. WASHINGTON, vol. 85, August 1988 (1988-08), page 5536-5540 XP002910776 ISSN: 0027-8424 the whole document	1

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/SG 99/00016

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9712512 A	10-04-1997	US 5846797 A AU 7257996 A CN 1198655 A EP 0910239 A	08-12-1998 28-04-1997 11-11-1998 28-04-1999
WO 9815622 A	16-04-1998	AU 4981297 A	05-05-1998
EP 0270355 A	08-06-1988	US 5004863 A AT 102999 T DE 3789359 D DE 3789359 T ES 2052582 T IN 168950 A US 5608142 A US 5159135 A	02-04-1991 15-04-1994 21-04-1994 06-10-1994 16-07-1994 20-07-1991 04-03-1997 27-10-1992
EP 0317511 A	24-05-1989	AU 616444 B AU 2568188 A JP 1160480 A RU 2024613 C	31-10-1991 18-05-1989 23-06-1989 15-12-1994
WO 8905344 A	15-06-1989	AT 178353 T AU 632038 B AU 2926689 A AU 668915 B AU 3528493 A AU 708250 B AU 6424796 A CA 1337406 A CA 1335799 A CN 1034298 A,B CN 1070309 A,B CN 1070310 A,B DE 3856319 D DE 3856319 T EP 0344302 A EP 0899341 A ES 2016428 A GR 88100761 A,B IL 88266 A IL 104845 A JP 7000065 A JP 8004434 B JP 8022196 B JP 2502253 T KR 9710757 B PT 89034 A,B US 5856177 A US 5834292 A US 5859321 A US 5874662 A US 5695999 A US 5583036 A US 5244802 A	15-04-1999 17-12-1992 05-07-1989 23-05-1996 20-05-1993 29-07-1999 21-11-1996 24-10-1995 06-06-1995 02-08-1989 31-03-1993 31-03-1993 06-05-1999 15-07-1999 06-12-1989 03-03-1999 01-11-1990 31-03-1994 10-03-1998 16-08-1998 06-01-1995 24-01-1996 06-03-1996 26-07-1990 30-06-1997 30-11-1989 05-01-1999 10-11-1998 12-01-1999 23-02-1999 09-12-1997 10-12-1996 14-09-1993
US 4672035 A	09-06-1987	NONE	